




種豬基因檢測技術： (一)緊迫與多產基因


廖仁寶副研究員
畜產試驗所遺傳育種組



豬染色體



www.animalgenome.org



Federico et al., 2004




大綱

- 豬遺傳背景
- 緊迫基因檢測技術
- 多產基因檢測技術
- 豬基因體研究現況
- 結論 Q&A




染色體組成



whyimportant.wordpress.com

DNA (去氧核醣核酸)




biotech.nstm.gov.tw/02/025.asp

緊迫基因 PSS

- 豬隻緊迫基因是一種**隱性遺傳**突變基因，與豬隻**暴斃**、肉豬屠體呈蒼白軟性**水樣肉**有關，對養豬業者造成經營上的極大損失。帶有此突變基因的豬容易在交配、運送屠宰的過程或受到驚嚇的狀況下呈現呼吸增快、肌肉僵直、心律不整、休克死亡等緊迫症狀。因為豬緊迫症狀所造成養豬業者在經濟上的損失，因此緊迫基因豬之篩除便成為一項重要的育種工作。帶有緊迫基因的豬通常對於麻醉藥劑的刺激特別敏感，而會引起與豬隻緊迫症狀相同的病症。早期在篩除緊迫基因豬是利用**鹵乙烷**測試法來檢驗，但是這種測試法對於受測仔豬是一種很強烈而且有可能致命的刺激。目前則是應用高效率、低成本及高準確性的分子生物技術檢測仔豬DNA以篩除緊迫基因。


豬第6號染色體



<http://www.marc.usda.gov/genome/htmls/LinkageMap.jsp>

緊迫基因

蒼白軟性水樣肉




鹵乙烷檢測





血液組成



tbmta.pixnet.net/blog/category/1626606



參考方法



Meat Science, Vol. 43, No. 2, 93-97, 1996
Copyright © 1996 Elsevier Science Ltd
Printed in Great Britain. All rights reserved
0309-1740/96 \$15.00 + 0.00
PII: S0309-1740(96)00006-X

Use of Mutagenically Separated PCR for the Detection of the Mutation Associated with Porcine Stress Syndrome

Andrew K. Lockley,* Jennifer S. Bruce, Stephen J. Franklin
& Ronald G. Bardsley

Department of Applied Biochemistry and Food Science, University of Nottingham, Sutton Bonington Campus, Loughborough, Leicestershire LE12 5RD, UK

(Received 14 November 1995; revised version received 10 January 1996;
accepted 12 January 1996)

ABSTRACT

A point mutation in the *Ryr1* gene encoding the ryanodine receptor in porcine skeletal muscle is associated with enhanced growth characteristics and leanness but also with porcine stress syndrome and pale, soft exudative meat in some animals. The current diagnostic test for the mutation is based on the polymerase chain reaction (PCR), followed by a restriction enzyme digestion step, prior to agarose gel electrophoresis. Using a technique known as mutagenically separated PCR (MS-PCR), a one-step procedure for the identification of the point mutation associated with porcine stress syndrome has been developed. This removes the requirement of the current PCR-based test for restriction enzyme digestion, is consequently quicker to perform, and may lend itself more readily to automation. DNA from blood samples from a series of animals were genotyped using both the conventional test and MS-PCR, and complete agreement between the two methods was obtained. Copyright © 1996 Elsevier Science Ltd



檢測方法

- 血液、精液、組織
- DNA萃取
- MSPCR
- 電泳分離
- 染色成像




PCR機器




welgene.pixnet.net

PCR原理




電泳槽



raphael426.pixnet.net

電泳圖



緊迫基因

出生年	品種數	場數	緊迫基因檢測				
			總頭數	BB	AB	AA	AA(%)
2014	10	18	1540	1	67	1472	95.6
2013	13	25	3013	2	85	2926	97.1
2012	12	33	3865	6	141	3718	96.2
2011	14	36	3845	4	208	3633	94.5
2010	12	39	3867	7	244	3616	93.5
2009	12	33	3404	10	240	3154	92.7
2008	8	31	2717	16	290	2411	88.7
2007	6	30	2973	8	332	2633	88.6
2006	6	27	3017	22	474	2521	83.6
2005	5	25	3272	44	617	2611	79.8

http://www.angrin.tli.gov.tw/pig_all.htm



藍瑞斯 Landrace

出生年	場數	緊迫基因檢測				
		總頭數	BB	AB	AA	AA(%)
2014	12	516	0	0	516	100.0
2013	17	986	0	3	983	99.7
2012	22	1079	1	6	1072	99.4
2011	22	1090	0	12	1078	98.9
2010	24	1264	0	23	1241	98.2
2009	21	1106	2	36	1068	96.6
2008	20	880	4	60	816	92.7
2007	23	987	0	21	966	97.9
2006	21	774	0	15	759	98.1
2005	21	853	1	16	836	98.0

http://www.angrin.tlri.gov.tw/pig_all.htm

約克夏 Yorkshire

出生年	場數	緊迫基因檢測				
		總頭數	BB	AB	AA	AA(%)
2014	9	149	0	2	147	98.7
2013	10	257	0	4	253	98.4
2012	14	330	0	3	327	99.1
2011	18	450	0	5	445	98.9
2010	15	418	0	3	415	99.3
2009	19	379	0	2	377	99.5
2008	16	334	0	2	332	99.4
2007	13	256	0	1	255	99.6
2006	14	204	0	9	195	95.6
2005	16	208	0	1	207	99.5

http://www.angrin.tlri.gov.tw/pig_all.htm



杜洛克 Duroc

出生年	場數	緊迫基因檢測				
		總頭數	BB	AB	AA	AA(%)
2014	15	639	1	46	592	92.6
2013	23	1356	2	64	1290	95.1
2012	25	1741	5	125	1611	92.5
2011	25	1860	4	167	1689	90.8
2010	29	2016	7	205	1804	89.5
2009	27	1797	8	197	1592	88.6
2008	28	1437	10	225	1202	83.6
2007	25	1713	8	310	1395	81.4
2006	23	2020	22	449	1549	76.7
2005	23	2184	43	598	1543	70.7

http://www.angrin.tlri.gov.tw/pig_all.htm



ESR多產基因標記

- 母豬的每胎產仔數是高繁殖性能品系培育的一項重要經濟性狀指標。研究發現，ESR遺傳標記與豬隻繁殖性能有關。結果顯示，具「多產基因純合型」的母豬，其第一胎產仔數平均每窩可多生0.39頭活仔豬，爾後胎次平均每窩亦可多生0.31頭。以不牽涉國際專利權且效率高、低成本之DNA檢測法(MS-PCR)，於仔豬階段進行早期選種計畫，不僅可減少所需更新候補種豬的飼養頭數，且可降低單位產能之生產成本，提升產業的競爭力。

參考資料

Effect of the Estrogen Receptor Locus on Reproduction and Production Traits in Four Commercial Pig Lines¹

T. H. Short*, M. F. Rothschild†, O. I. Southwood‡, D. G. McLaren*, A. de Vries‡, H. van der Steen‡, G. R. Eckard‡, C. K. Tuggle‡, J. Helm‡, D. A. Vaske‡, A. J. Mileham§, and G. S. Plastow§


*PIC USA, Franklin, KY 42135; †Department of Animal Science, Iowa State University, Ames 50011; ‡PIC Europe, Fyfield Wick, Abingdon, Oxfordshire, UK; §PIC Group, Fyfield Wick, Abingdon, Oxfordshire, UK; and §Marshfield Clinics, Marshfield, WI 54449

ABSTRACT: We investigated the effect of the estrogen receptor (ESR) gene on growth and reproductive traits in four Large White-based commercial pig lines. A total of 9,015 litter records from 4,262 sows genotyped at the ESR locus were analyzed to determine whether ESR influenced total number born (TNB) or number born alive (NBA). Teat number (TN), test ADG, ADFI, feed/gain ratio (F/G), and ultrasonic backfat (BF) were also analyzed to determine effects of ESR. The TNB and NBA were increased per favorable allele of ESR ($P < .01$) with additive effects of .42 (.31) and .39 (.31) pigs/litter in the first parity (later parities), respectively.


Key Words: Estrogen Receptors, Reproduction, Growth, Genetic Markers, Major Genes, Pleiotropy

J. Anim. Sci. 1997. 75:3138–3142

豬第1號染色體



ESR電泳圖



豬 ESR 基因型檢測

Panel I: MS-PCR Panel II: PCR/*Pvu*II

NN : 1、3、4、8

MN : 2、9、10

MM : 5、6、7

多產基因 ESR

年代	品種數	場數	多產基因檢測						
			總頭數	NN	MN	MM	MM(%)	M-	M-(%)
2014	10	18	1481	1164	198	119	8.0	436	14.7
2013	13	25	2948	2338	363	247	8.4	857	14.5
2012	12	33	3751	2899	499	353	9.4	1205	16.1
2011	14	36	3706	2997	413	296	8.0	1005	13.6
2010	12	38	3833	3274	386	173	4.5	732	9.5
2009	12	32	3222	2779	300	143	4.4	586	9.1
2008	8	31	2640	2274	247	119	4.5	485	9.2
2007	6	27	2275	2022	176	77	3.4	330	7.3
2006	6	25	2274	2147	87	40	1.8	167	3.7
2005	5	23	2538	2397	99	42	1.7	183	3.6

http://www.angrin.tlri.gov.tw/pig_all.htm

● ● ●

藍瑞斯 Landrace

出生年	場數	多產基因檢測						
		總頭數	NN	MN	MM	MM(%)	M-	M-(%)
2014	12	516	437	63	16	3.1	95	9.2
2013	17	985	841	114	30	3.0	174	8.8
2012	22	1072	912	149	11	1.0	171	8.0
2011	22	1078	957	119	2	0.2	123	5.7
2010	24	1242	1057	172	13	1.0	198	8.0
2009	21	1036	920	110	6	0.6	122	5.9
2008	20	816	747	69	0	0.0	69	4.2
2007	20	716	647	68	1	0.1	70	4.9
2006	17	543	515	26	2	0.4	30	2.8
2005	20	605	573	29	3	0.5	35	2.9

http://www.angrin.tlri.gov.tw/pig_all.htm

● ● ●

約克夏 Yorkshire

出生年	場數	多產基因檢測						
		總頭數	NN	MN	MM	MM(%)	M-	M-(%)
2014	9	149	13	81	55	36.9	191	64.1
2013	10	255	37	114	104	40.8	322	63.1
2012	14	328	66	146	116	35.4	378	57.6
2011	18	446	115	190	141	31.6	472	52.9
2010	15	416	97	184	135	32.5	454	54.6
2009	19	362	93	157	112	30.9	381	52.6
2008	16	332	68	170	94	28.3	358	53.9
2007	12	213	44	106	63	29.6	232	54.5
2006	12	134	44	60	30	22.4	120	44.8
2005	13	149	47	65	37	24.8	139	46.6


http://www.angrin.tlri.gov.tw/pig_all.htm

● ● ●

杜洛克 Duroc

出生年	場數	多產基因檢測						
		總頭數	NN	MN	MM	MM(%)	M-	M-(%)
2014	15	598	593	5	0	0.0	5	0.4
2013	23	1303	1302	1	0	0.0	1	0.0
2012	25	1637	1637	0	0	0.0	0	0.0
2011	25	1746	1745	1	0	0.0	1	0.0
2010	28	2006	2006	0	0	0.0	0	0.0
2009	26	1702	1697	3	2	0.1	7	0.2
2008	28	1428	1427	1	0	0.0	1	0.0
2007	23	1329	1328	1	0	0.0	1	0.0
2006	23	1578	1576	1	1	0.1	3	0.1
2005	22	1757	1752	5	0	0.0	5	0.1

http://www.angrin.tlri.gov.tw/pig_all.htm





豬基因體研究現況

Assembly:	Sscrofa10.2, Aug 2011
Database version:	79.102
Base Pairs (鹼基配對)	3,024,658,544
Golden Path Length:	2,808,525,991
Genebuild by:	Ensembl
Genebuild method:	Full genebuild
Genebuild started:	Sep 2011
Genebuild released:	May 2012
Genebuild last updated/patched:	Feb 2014
Coding genes: (蛋白質編碼基因)	21,630
Non coding genes:	3,124
Pseudogenes:	568
Gene transcripts:	30,585

www.ensembl.org/Sus_scrofa/Location/Genome



結論

- 利用先進之分子檢測技術(molecular biotechnology) 可有效地提升種豬(breeding pig)之選育(selection)成效。
- 種豬之基因檢測與選育，需政府機構(government)與民間業者(farmer)之相互合作(cooperation)，始克其功。



豬經濟性狀基因座資料庫 (Pig QTLdb)

- 12,618個數量性狀基因座 (QTLs)
- 656種不同性狀 (肉與屠體品質、健康、外表、產量、繁殖)

染色體	QTL	染色體	QTL	染色體	QTL
1	1933	8	714	15	500
2	1563	9	571	16	278
3	408	10	255	17	249
4	951	11	222	18	243
5	511	12	457	X	378
6	1199	13	529	Y	1
7	1451	14	611		

www.animalgenome.org/cgi-bin/QTLdb/SS/index



謝謝聆聽

敬請指教Q&A